

\mathcal{P} -kernel normal systems for \mathcal{P} -inversive semigroups

Hengwu Zheng

Received: 20 January 2010 / Accepted: 29 June 2010 / Published online: 9 July 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract As a generalization of Preston's kernel normal systems, \mathcal{P} -kernel normal systems for \mathcal{P} -inversive semigroups are introduced, and strongly regular \mathcal{P} -congruences on \mathcal{P} -inversive semigroups in terms of their \mathcal{P} -kernel normal systems are characterized. These results generalize the corresponding results for \mathcal{P} -regular semigroups and \mathcal{P} -inversive semigroups.

Keywords E-inversive semigroup · \mathcal{P} -inversive semigroup · Strongly regular \mathcal{P} -congruence · \mathcal{P} -kernel normal system

1 Introduction

For standard terminology and notation in semigroup theory see Howie [11]. As usual, $E(S)$ is the set of idempotents of a semigroup S , $V(a) = \{a' \in S : aa'a = a, a'aa' = a'\}$ is the set of all *inverses* of $a \in S$.

A semigroup S is *E-inversive* if for every $a \in S$ there exists $x \in S$ such that ax is idempotent. This concept was introduced by Thierrin [25]. A semigroup is an *E-semigroup* if its idempotents form a subsemigroup. Basic properties of E-inversive semigroups were given by Blyth and Almeida Santos [1], Catino and Miccoli [2], Hayes [9, 10], Mitsch and Petrich [18] and Mitsch [15–17]. From Lemma 3.1 in [15], a semigroup S is E-inversive if and only if

$$(\forall a \in S) \quad W(a) = \{a' \in S : a' = a'aa'\} \neq \emptyset.$$

The elements of $W(a)$ are called *weak inverses* of a .

Communicated by Thomas E. Hall.

H. Zheng (✉)

School of Mathematical Sciences, Qufu Normal University, Qufu, Shandong 273165, P.R. China
e-mail: hwzheng163@163.com

Preston [21] introduced kernel normal systems to characterize congruences on inverse semigroups. Meakin [14] gave a generalization of kernel normal systems for orthodox semigroups. Imaoka [12] obtained a generalization of Preston's kernel normal systems for regular $*$ -semigroups. A semigroup S is a *regular $*$ -semigroup* if S with a unary operation $*$ satisfies $(a^*)^* = a$, $(ab)^* = b^*a^*$ and $aa^*a = a$ for any $a, b \in S$ (see [12, 19]). We have seen that orthodox semigroups and regular $*$ -semigroups are within the class of \mathcal{P} -regular semigroups. A *\mathcal{P} -regular semigroup $S(P)$* is a regular semigroup S whose set of idempotents $E(S)$ includes a subset P with properties $P^2 \subseteq E(S)$, $pPp \subseteq P$ for all $p \in P$, and $\{a' \in V(a) : aP^1a' \cup a'P^1a \subseteq P\} \neq \emptyset$ for all $a \in S$, where $P^1 = P \cup \{1\}$ (see [28, 29]). Sen [23] gave a characterization of \mathcal{P} -congruences (the terminology a *\mathcal{P} -congruence* for a usual congruence on a \mathcal{P} -regular semigroup [28]) on \mathcal{P} -regular semigroups in terms of their \mathcal{P} -kernel normal systems.

The existence of the least group congruence on an E-inversive semigroup was noted by Hall and Munn [8]. An explicit description of this congruence was given by Mitsch [15]. The least group congruence on an E-inversive E-semigroup was characterized by Reither [22]. Further characterizations of group congruences on an E-inversive semigroup has been obtained by the author [30]. An alternative description of this congruence on an E-inversive E-semigroup was given by Weipoltshammer [27]. Blyth and Almeida Santos [1] gave an alternative description of group congruences on an E-inversive semigroup.

Weipoltshammer [27] described some congruences on E-inversive E-semigroups. References [4, 5] and [3] refer to work by Fan and Chen in which \mathcal{P} -inversive semigroups were first studied (using the so called \mathcal{P} -kernel normal systems) and the regular \mathcal{P} -congruences on \mathcal{P} -inversive semigroups. A *\mathcal{P} -inversive semigroup* is an E-inversive semigroup whose set of idempotents includes a subset P with properties that make it suitable in a “kernel normal systems” that generalizes the systems of Preston [21] and Meakin [14] in their respective congruence theories for inverse and orthodox semigroups.

Definition 1.1 (see [3–5]) An E-inversive semigroup S is called a *\mathcal{P} -inversive semigroup*, if there exists a nonempty subset P of $E(S)$ such that

- (1) $P^2 \subseteq E(S)$;
- (2) $(\forall p \in P) pPp \subseteq P$;
- (3) $(\forall a \in S) W_P(a) = \{a' \in W(a) : aP^1a' \cup a'P^1a \subseteq P\} \neq \emptyset$, where $P^1 = P \cup \{1\}$.

The subset P of $E(S)$ satisfying (1)–(3) above is called a *characteristic set* (*C-set*, for short) of S , each element in $W_P(a)$ is called a *weak \mathcal{P} -inverse* of a . Clearly, for any $p \in P$, we have $p \in W_P(p)$. Because of the central role of P in S we denote a \mathcal{P} -inversive semigroup S with the *C-set* P by $S(P)$. Throughout this paper, $S(P)$ is always an arbitrary \mathcal{P} -inversive semigroup.

From Definition 1.1 it follows that all \mathcal{P} -regular semigroups and E-inversive E-semigroups are within the class of \mathcal{P} -inversive semigroups. We have seen that there exist \mathcal{P} -inversive semigroups which are neither \mathcal{P} -regular nor E-inversive E-semigroups (see [4] and [7]).

A *regular \mathcal{P} -congruence* ρ on $S(P)$ is a congruence ρ with property $\rho a a a' a$ for all $a \in S(P)$ and all $a' \in W_P(a)$ in the sense of [3].

In [7], the strong \mathcal{P} -congruences and some sublattices of the strong \mathcal{P} -congruence lattice on a \mathcal{P} -inversive semigroup were studied. A *strong \mathcal{P} -congruence* ρ on $S(P)$ is a congruence ρ with properties for all $a, b \in S(P)$, $a\rho b$ implies $a'\rho b'$ for all $a' \in W_P(a)$, $b' \in W_P(b)$, and $apaa'a$ for all $a \in S(P)$ and all $a' \in W_P(a)$ in the sense of [4]. Siripitukdet and Sattayaporn [24] characterized the maximum idempotent separating congruence on an E-inversive E-semigroup.

The article [13] by Lou, Fan and Li is a major comprehensive paper on this topic. In it, any regular congruence on an arbitrary E-inversive semigroup is uniquely specified in terms of its trace and kernel (generalizing results for regular semigroups by Feigenbaum [6], Trotter [26] and especially by Pastijn and Petrich [20]). In the final section of [13], the regular congruences on E-inversive semigroups are characterized in terms of kernel normal systems.

The purpose of this paper is to generalize Preston's kernel normal systems to \mathcal{P} -inversive semigroups, and to give a description of strongly regular \mathcal{P} -congruences on \mathcal{P} -inversive semigroups in terms of their \mathcal{P} -kernel normal systems. The type of technique used here is basically the one used in [23]. These results generalize the corresponding results for \mathcal{P} -regular semigroups [23] and \mathcal{P} -inversive semigroups [3, 5].

By using the weak inverses in semigroups, Lou etc. [13] described regular congruences on E-inversive semigroups in terms of their kernel normal systems. In this paper we show that weak \mathcal{P} -inverses indeed can replace weak inverses in a congruence theory for \mathcal{P} -inversive semigroups, in a neater manner than that obtained in [13].

Recall that a congruence ρ on a semigroup S is said to be *regular* if S/ρ is a regular semigroup. Obviously, a congruence satisfying the property

$$(\forall a \in S) (\exists a' \in W_P(a)) \quad apaa'a \quad (\text{P})$$

on $S(P)$ is regular. In general, regular congruences on $S(P)$ may not satisfy the property (P). For example, let $S = (\mathbb{N}, \cdot)$ be the multiplicative semigroup of all non-negative integers and ρ be the congruence determined by the partition:

$$\{0\}, \{m \in \mathbb{N} : m > 0\}.$$

Then S is an E-inversive semigroup with $E(S) = \{0, 1\}$, and $S/\rho = \{\bar{0}, \bar{1}\}$ is a band. Let $P = \{0\}$. Certainly, $S(P)$ is \mathcal{P} -inversive, and ρ is regular. But there is no weak \mathcal{P} -inverse x of 1 such that $1\rho 1x1$. In this paper we shall be interested in the regular congruences which possesses the property (P) and we call such congruences *strongly regular*. That is,

Definition 1.2 A congruence ρ on $S(P)$ is called a *strongly regular \mathcal{P} -congruence*, if it satisfies

$$(\forall a \in S) (\exists a' \in W_P(a)) \quad apaa'a.$$

Example 1.3 Let S be the semigroup with Cayley table [27, Example 5.1]):

S	a	b	c	d	e	f	g
a	e	g	a	e	e	g	e
b	c	d	e	d	e	e	f
c	e	f	c	e	e	f	e
d	e	d	e	d	e	e	e
e	e	e	e	e	e	e	e
f	c	e	e	e	e	e	f
g	a	e	e	e	e	e	g

Then S is an E-inversive E-semigroup with $E(S) = \{c, d, e, g\}$. Let $P = E(S)$. Then $S(P)$ is \mathcal{P} -inverse. Let ρ be the congruence determined by the partition $\{a\}, \{b, f\}, \{c\}, \{d, e\}, \{g\}$. Notice that b is the only non-regular element of S , $a \in W_P(b)$, and $bab = f\rho b$, we have that ρ is a strongly regular \mathcal{P} -congruence on $S(P)$. Now $d \in W_P(b)$ and $bdb = d$, but d and b are not ρ -related. This implies that ρ is not a regular \mathcal{P} -congruence on $S(P)$. Certainly, ρ is not a strong \mathcal{P} -congruence on $S(P)$.

The previous example illustrates that the concept of strong regular \mathcal{P} -congruences on \mathcal{P} -inverse semigroups is a generalization of the concept of strong \mathcal{P} -congruences and regular \mathcal{P} -congruences on this class of semigroups.

2 \mathcal{P} -kernel normal systems

The set $\mathcal{B} = \{B_i : i \in I\}$ of subsemigroups of $S(P)$ is said to be a \mathcal{P} -kernel normal system for $S(P)$ if the following conditions hold:

- (K1) $B_i \cap B_j = \emptyset$ if $i \neq j \in I$.
- (K2) Each B_i contains an element of P and each element of P is contained in some B_i .
- (K3) For any $a \in S(P)$ there exists $a' \in W_P(a)$ such that

$$(\forall x, y \in S^1) (\forall B \in \mathcal{B}) xay \in B \Leftrightarrow xaa'ay \in B.$$

(K4)

$$(\forall x, y \in S^1) (\forall i_1, i_2, \dots, i_n, j \in I) \\ xB_{i_1}B_{i_2} \cdots B_{i_n}y \cap B_j \neq \emptyset \Rightarrow xB_{i_1}B_{i_2} \cdots B_{i_n}y \subseteq B_j.$$

The weak \mathcal{P} -inverse a' of a satisfying (K3) above is called a weak \mathcal{B} -inverse of a , and $W_B(a)$ denotes the set of weak \mathcal{B} -inverses of a . Now $W_B(a) \neq \emptyset$ for any $a \in S(P)$.

Let ρ be a strongly regular \mathcal{P} -congruence on $S(P)$. The set $\{p\rho : p \in P\}$ is denoted by \mathcal{B}_ρ . The following lemma also gives the examples of \mathcal{P} -kernel normal systems for $S(P)$.

Lemma 2.1 *If ρ is a strongly regular \mathcal{P} -congruence on $S(P)$, then \mathcal{B}_ρ is a \mathcal{P} -kernel normal system for $S(P)$.*

Proof (K1) and (K2) are clear.

(K3) Since ρ is a strongly regular \mathcal{P} -congruence, for any $a \in S(P)$ there exists $a' \in W_P(a)$ such that $apaa'a$. Obviously, $a' \in W_B(a)$.

(K4) Let $b = xb_{i_1}b_{i_2} \cdots b_{i_n}y \in xB_{i_1}B_{i_2} \cdots B_{i_n}y \cap B_j$, where $b_{i_t} \in B_{i_t}$, $t = 1, 2, \dots, n$, and let $B_j = p\rho$, $p \in P$. Then $b\rho p$. For any $c = xc_{i_1}c_{i_2} \cdots c_{i_n}y \in xB_{i_1}B_{i_2} \cdots B_{i_n}y$, where $c_{i_t} \in B_{i_t}$, $t = 1, 2, \dots, n$, we have that $b_{i_t}, c_{i_t} \in B_{i_t}$ implies $b_{i_t}\rho = c_{i_t}\rho$, and so that $c\rho = x\rho c_{i_1}\rho c_{i_2}\rho \cdots c_{i_n}\rho y\rho = x\rho b_{i_1}\rho b_{i_2}\rho \cdots b_{i_n}\rho y\rho = b\rho = p\rho$. Hence $c \in B_j$. \square

We have the following results on \mathcal{P} -kernel normal systems which will be useful in the sequel.

Lemma 2.2 *Let $\mathcal{B} = \{B_i : i \in I\}$ be a \mathcal{P} -kernel normal system for $S(P)$.*

- (1) *If $a' \in W_B(a)$, $b' \in W_B(b)$, then $aa'a \in W_B(a')$, $b'a' \in W_B(ab)$.*
- (2) *Let $x, y \in S^1$, $i_1, i_2, \dots, i_n \in I$. If $xB_{i_1}B_{i_2} \cdots B_{i_n}y \cap P \neq \emptyset$, then there exists $j \in I$ such that $xB_{i_1}B_{i_2} \cdots B_{i_n}y \subseteq B_j$.*
- (3) *For $a \in S(P)$, $a' \in W_B(a)$ and $i \in I$, there exist $j, k \in I$ such that $aB_i a' \subseteq B_j$, and $a'B_i a \subseteq B_k$.*
- (4) *If $a \in B_i$, $a' \in W_B(a)$ then there exist $j, k \in I$ such that $a'B_i \subseteq B_j$ and $B_i a' \subseteq B_k$.*
- (5) *If $a, ab \in B_i$ and $ab', bb' \in B_j$ for some $b' \in W_B(b)$, $i, j \in I$, then $b \in B_i$.*
- (6) *If $a, ba \in B_i$ and $b'a, b'b \in B_j$ for some $b' \in W_B(b)$, $i, j \in I$, then $b \in B_i$.*

Proof (1) Let $a' \in W_B(a)$, $b' \in W_B(b)$. It is easy to see that $aa'a \in V(a')$ and $aa'a \in W_P(a')$. Thus $aa'a \in W_B(a')$.

Now $a' \in W_P(a)$, $b' \in W_P(b)$. It follows from [4, Lemma 1.4] that $b'a' \in W_P(ab)$. Let $x, y \in S^1$. Then

$$\begin{aligned} xabb'a'aby \in B_i &\Leftrightarrow xaa'abb'a'abb'by \in B_i \quad (\text{since } a' \in W_B(a) \text{ and } b' \in W_B(b)) \\ &\Leftrightarrow xa(a'abb')^2by \in B_i \\ &\Leftrightarrow xaa'abb'by \in B_i \quad (\text{since } a'abb' \in P^2 \subseteq E(S)) \\ &\Leftrightarrow xaby \in B_i \quad (\text{since } a' \in W_B(a) \text{ and } b' \in W_B(b)). \end{aligned}$$

Hence $b'a' \in W_B(ab)$.

(2) By (K2), each element of P is contained in some B_j . The result follows from (K4).

(3) Suppose that B_i contains some element $p \in P$. Then $apa' \in P$. Hence $aB_i a' \cap P \neq \emptyset$. Now it follows from (2) that there exists $j \in I$ such that $aB_i a' \subseteq B_j$. Similarly, we can show the other result.

(4) Since $a \in B_i$ and $a' \in W_B(a)$, $a'a \in a'B_i \cap P$. It follows from (2) that there exists $B_j \in \mathcal{B}$ such that $a'B_i \subseteq B_j$. Similarly, we can prove the other result.

(5) Since $ab \in B_i, abb'b \in B_i$. Now $b'b \in P$, by (K2), there exists $B_k \in \mathcal{B}$ such that $b'b \in B_k$. Hence $ab'b, abb'b \in B_i B_k$, and so $B_i B_k \cap B_i \neq \emptyset$. By (K4), $B_i B_k \subseteq B_i$, so that $ab'b \in B_i$. Now $ab'b \in B_j b$. Thus $B_j b \cap B_i \neq \emptyset$. By (K4), $B_j b \subseteq B_i$. Since $bb' \in B_j, bb'b \in B_i$, that is, $b \in B_i$.

(6) Similar to the proof of (5), we can obtain (6). \square

We now introduce the following notation. Let $\mathcal{B} = \{B_i : i \in I\}$ be a \mathcal{P} -kernel normal system for $S(P)$ and $a, b \in S(P)$. By $a \sim b$ we mean that a and b are contained in a same B_i . Define a relation $\rho_{\mathcal{B}}$ on $S(P)$ by

$$a\rho_{\mathcal{B}}b \Leftrightarrow (\exists a' \in W_B(a))(\exists b' \in W_B(b))ba' \sim aa' \text{ \& \& } b'a \sim b'b.$$

Clearly, $\rho_{\mathcal{B}}$ is a reflexive relation on $S(P)$.

In order to obtain the main result in this paper, we shall set up a series of lemmas.

Lemma 2.3 $\rho_{\mathcal{B}}$ is a symmetric relation on $S(P)$.

Proof Let $a\rho_{\mathcal{B}}b$. Then there exist $a' \in W_B(a), b' \in W_B(b)$ such that $ba' \sim aa'$ and $b'a \sim b'b$. Then there exist $B_{i_1}, B_{i_2} \in \mathcal{B}$ such that $ba', aa' \in B_{i_1}$ and $b'a, b'b \in B_{i_2}$. Let $a^+ \in W_B(a)$ and $b^+ \in W_B(b)$. Now by Lemma 2.2(1), $aa'ab^+ \in W_B(ba')$. Then by Lemma 2.2(4), $B_{i_1}(aa'ab^+) \subseteq B_{i_3}$ for some $i_3 \in I$. Now $aa'ab^+ = (aa')aa'ab^+ \in B_{i_1}(aa'ab^+) \subseteq B_{i_3}$. By (K3), $ab^+ \in B_{i_3}$. Also, $ba'ab^+ = (ba')(aa'a)b^+ \in B_{i_3}$. Then by $b' \in W_B(b), b(b'b)(a'a)b^+ \in B_{i_3}$. Hence $bB_{i_2}B_{i_4}b^+ \cap B_{i_3} \neq \emptyset$, where $a'a \in B_{i_4}$. By (K4), $bB_{i_2}B_{i_4}b^+ \subseteq B_{i_3}$. Hence $bb'aa'ab^+ \in B_{i_3}$, that is, $bb'ab^+ \in B_{i_3}$. Hence $ab^+, bb'ab^+ \in B_{i_3}$. Now $b'a \sim b'b$ implies that $b(b'b)b^+ \sim b(b'a)b^+$ by Lemma 2.2(3). Therefore $ab^+ \sim bb'ab^+ \sim bb'bb^+ \sim bb^+$, and so $ab^+ \sim bb^+$. Similarly, we can prove that $a^+b \sim a^+a$. Hence $b\rho_{\mathcal{B}}a$, and so that $\rho_{\mathcal{B}}$ is symmetric. \square

Remark In the proof of Lemma 2.3, we also prove that if there exist $a' \in W_B(a), b' \in W_B(b)$ such that $ba' \sim aa'$ and $b'a \sim b'b$, then $ab^+ \sim bb^+$ and $a^+b \sim a^+a$ for all $a^+ \in W_B(a), b^+ \in W_B(b)$. In the similar way, we can prove that if there exist $a^+ \in W_B(a), b^+ \in W_B(b)$ such that $ab^+ \sim bb^+$ and $a^+b \sim a^+a$, then $ba' \sim aa'$ and $b'a \sim b'b$ for all $a' \in W_B(a), b' \in W_B(b)$. In the previous definition of $\rho_{\mathcal{B}}$ we can substitute “there exists” by “for all”.

Lemma 2.4 If $(a, b), (b, c) \in \rho_{\mathcal{B}}$, then $a(c'c)a' \sim aa', c(a'a)c' \sim cc', a'(cc')a \sim a'a, b'(aa')b \sim b'b$ for all $a' \in W_B(a), b' \in W_B(b), c' \in W_B(c)$.

Proof Since $(a, b), (b, c) \in \rho_{\mathcal{B}}, ba' \sim aa', b'a \sim b'b, a'b \sim a'a, ab' \sim bb', cb' \sim bb', c'b \sim c'c, b'c \sim b'b, bc' \sim cc'$ for all $a' \in W_B(a), b' \in W_B(b)$ and $c' \in W_B(c)$ (from the remark after Lemma 2.3). So there exist $B_{i_1}, B_{i_2}, B_{i_3}, B_{i_4}, B_{i_5}, B_{i_6}, B_{i_7}$ and B_{i_8} in \mathcal{B} such that $ba', aa' \in B_{i_1}, b'a, b'b \in B_{i_2}, a'b, a'a \in B_{i_3}, ab', bb' \in B_{i_4}, cb', bb' \in B_{i_5}, c'b, c'c \in B_{i_6}, b'c, b'b \in B_{i_7}$ and $bc', cc' \in B_{i_8}$. From (K1), $B_{i_2} = B_{i_7}$ and $B_{i_4} = B_{i_5}$.

We first show that $a(c'c)a' \sim aa'$. Now $a(a'b)(b'b)(c'c)a'$, $a(a'b)(b'c)(c'c)a'$, $a(a'a)(b'c)(c'b)a' \in aB_{i_3}B_{i_2}B_{i_6}a'$. Since $a(b'c)(c'b)a' \in P$, by (K2), there exists $i_9 \in I$ such that $a(b'c)(c'b)a' \in B_{i_9}$, and so $a(a'a)(b'c)(c'b)a' \in B_{i_9}$ by (K3). Thus $aB_{i_3}B_{i_2}B_{i_6}a' \cap B_{i_9} \neq \emptyset$. Hence from (K4), $aB_{i_3}B_{i_2}B_{i_6}a' \subseteq B_{i_9}$, so that $a(a'b)(b'b)(c'c)a'$, $a(a'b)(b'c)(c'c)a' \in B_{i_9}$, that is, $a(a'b)(c'c)a'$, $a(a'b)(b'c)a' \in B_{i_9}$.

Now $a(a'a)(c'c)a'$, $a(a'b)(c'c)a' \in aB_{i_3}B_{i_6}a'$. Since $a(c'c)a' \in P$, there exists $i_{10} \in I$ such that $a(c'c)a' \in B_{i_{10}}$. Then by (K3), $a(a'a)(c'c)a' \in B_{i_{10}}$. Hence $aB_{i_3}B_{i_6}a' \cap B_{i_{10}} \neq \emptyset$. Then by (K4), $aB_{i_3}B_{i_6}a' \subseteq B_{i_{10}}$. Thus $a(a'b)(c'c)a' \in B_{i_{10}}$. By (K1), $B_{i_{10}} = B_{i_9}$. Hence $a(a'a)(c'c)a' \in B_{i_9}$, and so $a(c'c)a' \in B_{i_9}$.

Now $a(a'b)(b'b)a'$, $a(a'b)(b'c)a'$, $a(a'a)(b'b)a' \in aB_{i_3}B_{i_7}a'$. Note $a(b'b)a' \in P$, there exists $i_{11} \in I$ such that $a(b'b)a' \in B_{i_{11}}$. By (K3), $a(a'a)(b'b)a' \in B_{i_{11}}$, and so $aB_{i_3}B_{i_7}a' \cap B_{i_{11}} \neq \emptyset$. Then by (K4), $aB_{i_3}B_{i_7}a' \subseteq B_{i_{11}}$. Hence $a(a'b)(b'c)a' \in B_{i_{11}}$. By (K1), $B_{i_{11}} = B_{i_9}$. Then $a(c'c)a' \sim a(a'b)(b'b)a' \sim a(a'b)a'$. Now $a'b \sim a'a$ implies (by Lemma 2.2(3)) that $a(a'b)a' \sim a(a'a)a' = aa'$. Hence $a(c'c)a' \sim aa'$.

Similarly, we can prove the other results. \square

Lemma 2.5 $\rho_{\mathcal{B}}$ is a transitive relation on $S(P)$.

Proof Here we use all the notation of Lemma 2.4. Let $(a, b) \in \rho_{\mathcal{B}}$ and $(b, c) \in \rho_{\mathcal{B}}$. Let $x = bc'cb'aa'$. Now $(cb')(aa')$, $(bb')(ba') \in B_{i_5}B_{i_1}$ and $(bb')(ba') \sim ba' \in B_{i_1}$. Hence $B_{i_5}B_{i_1} \cap B_{i_1} \neq \emptyset$. Then by (K4), $B_{i_5}B_{i_1} \subseteq B_{i_1}$. Hence $(cb')(aa') \in B_{i_1}$. This implies that $x = (bc')(cb'aa')$, $(cc')(cb'aa') \in B_{i_8}B_{i_1}$. But $cb'aa' \sim (cc')(cb'aa')$. Hence $B_{i_8}B_{i_1} \cap B_{i_1} \neq \emptyset$. Then by (K4), $B_{i_8}B_{i_1} \subseteq B_{i_1}$. Hence $x \in B_{i_1}$.

Let $y = ca'$. By Lemma 2.2(1), $aa'ac' \in W_B(y)$. Let $y' = aa'ac'$. Now $a'b \sim a'a$ implies (by Lemma 2.2(3)) $c(a'b)c' \sim c(a'a)c'$. From Lemma 2.4, $ca'ac' \sim cc'$. Hence $c(a'b)c'$, $cc' \in B_{i_8}$. Now $(cb')(aa')$, $(bb')(ba') \in B_{i_5}B_{i_1}$. But $(bb')(ba') \sim ba' \in B_{i_1}$. Hence $B_{i_5}B_{i_1} \cap B_{i_1} \neq \emptyset$, and so $(cb')(aa') \in B_{i_1}$. Then $(ca'bc')(cb')(aa')$, $(cc')(cb')(aa') \in B_{i_8}B_{i_1}$. But $(cc')(cb')(aa') \sim (cb')(aa') \in B_{i_1}$. Thus $B_{i_8}B_{i_1} \cap B_{i_1} \neq \emptyset$, so that $B_{i_8}B_{i_1} \subseteq B_{i_1}$. Hence

$$yx = ca'bc'cb'aa' \in B_{i_1}.$$

Now $x, yx \in B_{i_1}$ implies that

$$yx \sim x. \quad (2.1)$$

To show that $y'y \sim y'x$, we notice that

$$a(c'b)(c'c)(b'c)a', a(c'b)(c'c)(b'a)a', a(c'c)(c'b)(b'b)a' \in aB_{i_6}B_{i_6}B_{i_7}a'.$$

Here $a(c'b)(c'c)(b'c)a' = a(c'(b(c'c)b')c)a' \in P$. Hence by Lemma 2.2(2), there exists i_{12} such that $aB_{i_6}B_{i_6}B_{i_7}a' \subseteq B_{i_{12}}$. Hence $a(c'c)(c'b)(b'b)a' \in B_{i_{12}}$, that is, $ac'ba' \in B_{i_{12}}$.

Now $c'c \sim c'b$. Hence by Lemma 2.2(3), $a(c'c)a' \sim a(c'b)a'$. But $a(c'c)a'$, $aa' \in B_{i_1}$ by Lemma 2.4. Hence $a(c'b)a' \in B_{i_1}$. By (K1), $B_{i_1} = B_{i_{12}}$. Consequently, $ac'bc'cb'aa' \in B_{i_1}$, and so, $aa'ac'bc'cb'aa' \in B_{i_1}$. Hence $y'x, x \in B_{i_1}$. Again

$ac'ca' \sim aa'$. Then $y'y = aa'ac'ca' \sim ac'ca' \sim aa' \sim x$. Therefore

$$y'x \sim y'y. \quad (2.2)$$

From (2.1), (2.2), by Lemma 2.2(6), we have $y \sim x$. Then

$$ca' \sim x \sim aa'.$$

Next we show that $c'a \sim c'c$. Let $x_1 = c'cb'aa'b$, $y_1 = c'a$. Now $a'cc'c \in W_B(y_1)$. Let $y'_1 = a'cc'c$. From Lemma 2.4, $b'aa'b \sim b'b \in B_{i_7}$. Hence $(c'c)(b'aa'b)$, $(c'c)(b'a)$, $(c'b)(b'b) \in B_{i_6}B_{i_7}$. But $(c'b)(b'b) \sim c'b \in B_{i_6}$, and so $B_{i_6}B_{i_7} \cap B_{i_6} \neq \emptyset$. Then (K4) implies that $B_{i_6}B_{i_7} \subseteq B_{i_6}$, so that $x_1 = (cc')(b'aa'b) \in B_{i_6}$.

Now $x_1y_1 = (c'cb'a)(a'b)(c'a) = (c'c)(b'a)(a'bc'a)$. Here $a'(bc'a) \sim a'(cc'a)$ (by Lemma 2.2(3)) $\sim a'a$ (by Lemma 2.4) $\in B_{i_3}$. Notice that $(c'c)(b'a) \in B_{i_6}$ and so $x_1y_1 \in B_{i_6}B_{i_3}$. Also, $(c'c)(b'a)(a'a) \in B_{i_6}B_{i_3}$, and $(c'c)(b'a)(a'a) \sim (c'c)(b'a) \in B_{i_6}$. Hence $B_{i_6}B_{i_3} \subseteq B_{i_6}$, so that $x_1y_1 \in B_{i_6}$. Now $x_1, x_1y_1 \in B_{i_6}$ implies that

$$x_1y_1 \sim x_1. \quad (2.3)$$

To show that $x_1y'_1 \sim y_1y'_1$, notice that $x_1y'_1 = c'cb'aa'ba'cc'c$. Now $(cb')(aa')$, $(bb')(ba') \in B_{i_5}B_{i_1}$. But $(bb')(ba') \sim ba' \in B_{i_1}$. Hence $(cb')(aa')$, $ba' \in B_{i_1}$. So $cb'aa'ba' = (cb'aa')(ba') \in B_{i_1}B_{i_1} \subseteq B_{i_1}$ (since B_{i_1} is a subsemigroup). Then $cb'aa'ba' \sim aa'$. It follows from Lemma 2.2(3) that $c'(cb'aa'ba')c \sim c'(aa')c$. Hence $c'(cb'aa'ba')cc'c \sim c'(aa')cc'c$, that is,

$$x_1y'_1 \sim y_1y'_1. \quad (2.4)$$

From (2.3), (2.4), by Lemma 2.2(5), we have that $x_1 \sim y_1$, and so $c'c \sim x_1 \sim c'a$. Hence $(a, c) \in \rho_{\mathcal{B}}$. \square

Lemma 2.6 $\rho_{\mathcal{B}}$ is a strongly regular \mathcal{P} -congruence on $S(P)$.

Proof It follows from Lemma 2.3 and Lemma 2.5 that $\rho_{\mathcal{B}}$ is an equivalence relation on $S(P)$. Let us show that $\rho_{\mathcal{B}}$ is left compatible. Let $(a, b) \in \rho_{\mathcal{B}}$, $c \in S$, $a' \in W_B(a)$, $b' \in W_B(b)$ and $c' \in W_B(c)$. Then $ba' \sim aa'$, $b'a \sim b'b$, $a'b \sim a'a$, $ab' \sim bb'$. So there exist $B_{i_1}, B_{i_2}, B_{i_3}, B_{i_4} \in \mathcal{B}$ such that $ba', aa' \in B_{i_1}$, $b'a, b'b \in B_{i_2}$, $a'b, a'a \in B_{i_3}$, $ab', bb' \in B_{i_4}$. Now $ba' \sim aa'$ implies that $c(ba')c' \sim c(aa')c'$ (by Lemma 2.2(3)).

We show that $b'c'ca \sim b'c'cb$ as follows. Let $x = b'c'cba'a$, $y = b'c'ca$, $y' = a'c'cbb'b$. Then $y' \in W_B(y)$. Notice that $b'c'cb \in P$. Let $b'c'cb \in B_{i_5} \in \mathcal{B}$. Now $(b'c'cb)(b'a)$, $(b'c'cb)(b'b) \in B_{i_5}B_{i_2}$. Since $(b'c'cb)(b'b) \sim b'c'cb \in B_{i_5}$, $B_{i_5}B_{i_2} \cap B_{i_5} \neq \emptyset$. Hence by (K4), $B_{i_5}B_{i_2} \subseteq B_{i_5}$. Then $(b'c'cb)(b'a) \in B_{i_5}$. Now $(b'c'cb)(b'b)(a'a)$, $(b'c'cb)(b'a)(a'a) \in B_{i_5}B_{i_3}$. But $(b'c'cb)(b'a)(a'a) \sim (b'c'cb)(b'a) \in B_{i_5}$ shows that $B_{i_5}B_{i_3} \cap B_{i_5} \neq \emptyset$. Then $B_{i_5}B_{i_3} \subseteq B_{i_5}$ and $(b'c'cb)(b'b)(a'a) \in B_{i_5}$, that is, $x = b'c'cba'a \in B_{i_5}$.

Now $ab' \sim bb'$, and $b'c'c \in W_B(c'cb)$. Hence by Lemma 2.2(3), $b'(c'c(ab')c'c)b \sim b'(c'c(bb')c'c)b$. Since $b'c'cb \in P \subseteq E_S$, $b'(c'c(bb')c'c)b = (b'c'cb)^2 = b'c'cb$.

Now $b'c'cb \in B_{i_5}$, hence $b'(c'(ab')c'c)b \in B_{i_5}$. Then $yx = (b'c'ca)(b'c'cba'a) = (b'(c'cab'c'c)b)(a'a) \in B_{i_5}B_{i_3}$. But $B_{i_5}B_{i_3} \subseteq B_{i_5}$ and so $yx \in B_{i_5}$. Then

$$yx \sim x \sim b'c'cb. \quad (2.5)$$

To show that $y'x \sim y'y$, notice that $(c'c)(bb')(c'c) \in P$. Then there exists $B_{i_6} \in \mathcal{B}$ such that $(c'c)(bb')(c'c) \in B_{i_6}$. Now $y'x = a'c'cbb'bb'c'cba'a = a'(c'cbb'c'c)(ba'a)a$, $a'(c'cbb'c'c)(aa'a)a \in a'B_{i_6}B_{i_1}a$. Notice that $y'y = a'c'cbb'bb'c'ca = a'(c'cbb'c'c)a \in P$. Then there exists $B_{i_7} \in \mathcal{B}$ such that $y'y \in B_{i_7}$, and so $a'(c'cbb'c'c)(aa'a)a \in B_{i_7}$. Hence $a'B_{i_6}B_{i_1}a \subseteq B_{i_7}$. Then

$$y'x \sim y'y. \quad (2.6)$$

From (2.5), (2.6), by Lemma 2.2(6), $y \sim x \sim b'c'cb$, that is, $b'c'ca \sim b'c'cb$.

By Lemma 2.2(1), $a'c' \in W_B(ca)$, $b'c' \in W_B(cb)$. Hence $(ca, cb) \in \rho_{\mathcal{B}}$, and so $\rho_{\mathcal{B}}$ is left compatible. A similar argument shows that $\rho_{\mathcal{B}}$ is right compatible.

It is easy to verify that $a' \in W_B(aa'a)$ and $(a, aa'a) \in \rho_{\mathcal{B}}$. Hence $\rho_{\mathcal{B}}$ is a strongly regular \mathcal{P} -congruence on $S(P)$. \square

Lemma 2.7 *If $p \in B_i \cap P$, then $p\rho_{\mathcal{B}} = B_i$.*

Proof Let $a \in p\rho_{\mathcal{B}}$. Then $p\rho_{\mathcal{B}}a$. Since $p \in W_B(p)$, $ap \sim pp = p$ and $a'p \sim a'a$ for any $a' \in W_B(a)$. Hence by Lemma 2.2(6), $a \in B_i$.

Conversely, let $b \in B_i$. Since B_i is a subsemigroup, $pb \in B_i$. Then $pb \sim pp$. Let $b' \in W_B(b)$. By Lemma 2.2(4), there exists $B_j \in \mathcal{B}$ such that $B_ib' \subseteq B_j$. Hence $pb', bb' \in B_j$, and so $pb' \sim bb'$. By the definition of $\rho_{\mathcal{B}}$, $(b, p) \in \rho_{\mathcal{B}}$. Thus $b \in p\rho_{\mathcal{B}}$. \square

Lemma 2.8 *Let ρ and σ be two strongly regular \mathcal{P} -congruences on $S(P)$. Then $\rho = \sigma$ if and only if $\mathcal{B}_{\rho} = \mathcal{B}_{\sigma}$.*

Proof It suffices to show the “if” part. Let apb . Since σ is a strongly regular \mathcal{P} -congruence, there exists $a' \in W_P(a)$ such that $a\sigma aa'a$. Now $aa'\rho ba'$. Note $aa' \in P$. By the assumption, there exists $p \in P$ such that $aa'\rho = p\sigma$. Hence $aa'\rho = aa'\sigma$, and so $ba' \in aa'\sigma$, that is, $aa'\sigma ba'$. Similarly, we have that $b\sigma bb'b$ and $b'a\sigma b'b$ for some $b' \in W_P(b)$. Thus

$$a\sigma aa'a\sigma ba'a\sigma bb'ba'a\sigma bb'aa'a\sigma bb'a\sigma bb'b\sigma b.$$

Hence $a\sigma b$, and so $\rho \subseteq \sigma$. Similarly, we have $\sigma \subseteq \rho$. Thus $\rho = \sigma$. \square

The following theorem is the main result in this paper.

Theorem 2.9 *If \mathcal{B} is a \mathcal{P} -kernel normal system for $S(P)$, then $\rho_{\mathcal{B}}$ is a strongly regular \mathcal{P} -congruence on $S(P)$ and $\mathcal{B}_{\rho_{\mathcal{B}}} = \mathcal{B}$. Conversely, if ρ is a strongly regular \mathcal{P} -congruence on $S(P)$, then \mathcal{B}_{ρ} is a \mathcal{P} -kernel normal system for $S(P)$ and $\rho_{\mathcal{B}_{\rho}} = \rho$.*

Proof Direct part. Let $\mathcal{B} = \{B_i : i \in I\}$ be a \mathcal{P} -kernel normal system for $S(P)$. By Lemma 2.6, $\rho_{\mathcal{B}}$ is a strongly regular \mathcal{P} -congruence on $S(P)$.

Let $p \in P$. Then there exists $B_i \in \mathcal{B}$ such that $p \in B_i$. Hence by Lemma 2.7, $p\rho_{\mathcal{B}} = B_i$, and so $\mathcal{B}_{\rho_{\mathcal{B}}} \subseteq \mathcal{B}$. Conversely, suppose that $B_i \in \mathcal{B}$. Then B_i contains some element $p \in P$. By Lemma 2.7 again, $B_i = p\rho_{\mathcal{B}}$, so that $\mathcal{B}_{\rho_{\mathcal{B}}} = \mathcal{B}$ as required.

Converse part. Let ρ be a strongly regular \mathcal{P} -congruence on $S(P)$. Then it follows from Lemma 2.1 that \mathcal{B}_{ρ} is a \mathcal{P} -kernel normal system for $S(P)$. From the direct part of this theorem, we have that $\rho_{\mathcal{B}_{\rho}}$ is also a strongly regular \mathcal{P} -congruence on $S(P)$ and $\mathcal{B}_{\rho_{\mathcal{B}_{\rho}}} = \mathcal{B}_{\rho}$. Hence by Lemma 2.8, $\rho_{\mathcal{B}_{\rho}} = \rho$. \square

The set of all strongly regular \mathcal{P} -congruences on $S(P)$ and the set of all \mathcal{P} -kernel normal systems for $S(P)$ are denoted by $SRC_P(S)$ and $KNS_P(S)$, respectively.

Define a relation \leq on $KNS_P(S)$ by

$$\mathcal{A} \leq \mathcal{B} \Leftrightarrow (\forall A \in \mathcal{A}) (\exists B \in \mathcal{B}) A \subseteq B.$$

It is clear that \leq is a partial order on $KNS_P(S)$.

Now the next result follows immediately.

Corollary 2.10 *The mappings*

$$\begin{aligned} \varphi : \quad SRC_P(S) &\rightarrow KNS_P(S) \\ \rho &\mapsto \mathcal{B}_{\rho} \end{aligned}$$

and

$$\begin{aligned} \psi : \quad KNS_P(S) &\rightarrow SRC_P(S) \\ \mathcal{B} &\mapsto \rho_{\mathcal{B}} \end{aligned}$$

are mutually inverse order preserving bijections.

Acknowledgements The author would like to thank Professor Thomas E. Hall and the referee for their valuable suggestions and comments.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Blyth, T.S., Almeida Santos, M.H.: E-inversive Dubreil-Jacotin semigroups. *Semigroup Forum* **78**, 511–527 (2009)
2. Catino, F., Miccoli, M.M.: On semidirect products of semigroups. *Note Math.* **9**, 189–194 (1989)
3. Fan, X.K., Chen, Q.H.: Regular \mathcal{P} -congruences on \mathcal{P} -inversive semigroups. *Pure Appl. Math.* **19**, 374–378 (2003)
4. Fan, X.K., Chen, Q.H.: Strong \mathcal{P} -congruences on \mathcal{P} -inversive semigroups. *Adv. Math.* **33**, 434–440 (2004)
5. Fan, X.K., Chen, Q.H.: \mathcal{P} -kernel normal system of strong \mathcal{P} -congruences on \mathcal{P} -inversive semigroups. *Northeast. Math. J.* **21**, 397–403 (2005)

6. Feigenbaum, R.: Regular semigroup congruences. *Semigroup Forum* **17**, 373–377 (1979)
7. Gao, Z.H., Yu, B.J.: Sublattices of the lattices of strong \mathcal{P} -congruences on \mathcal{P} -inversive semigroups. *Semigroup Forum* **75**, 272–292 (2007)
8. Hall, T.E., Munn, W.D.: The hypercore of a semigroup. *Proc. Edinb. Math. Soc.* **28**, 107–122 (1985)
9. Hayes, A.D.: E^* -dense E -semigroups. *Semigroup Forum* **68**, 218–232 (2004)
10. Hayes, A.D.: Semigroups which have a minimum primitive inverse congruence. *Semigroup Forum* **65**, 386–404 (2002)
11. Howie, J.M.: *Fundamentals of Semigroup Theory*. Clarendon, Oxford (1995)
12. Imaoka, T.: $*$ -congruences on regular $*$ -semigroups. *Semigroup Forum* **23**, 321–326 (1981)
13. Luo, Y.F., Fan, X.K., Li, X.L.: Regular congruences on an E -inversive semigroup. *Semigroup Forum* **76**, 107–123 (2008)
14. Meakin, J.: Congruences on orthodox semigroups. *J. Aust. Math. Soc.* **12**, 323–341 (1971)
15. Mitsch, H.: Subdirect products of E -inversive semigroups. *J. Aust. Math. Soc.* **48**, 66–78 (1990)
16. Mitsch, H.: On the lemma of Lallement. *Commun. Algebra* **24**, 3085–3098 (1996)
17. Mitsch, H.: Introduction to E -inversive semigroups. In: Smith, P., Giraldes, E., Martins, P. (eds.) *Proc. Conf. Semigroups*. World Scientific, Singapore (2000)
18. Mitsch, H., Petrich, M.: Basic properties of E -inversive semigroups. *Commun. Algebra* **28**, 5169–5182 (2000)
19. Nordahl, T.E., Scheiblich, H.E.: Regular $*$ -semigroups. *Semigroup Forum* **28**, 369–377 (1978)
20. Pastijn, F., Perich, M.: Congruences on regular semigroups. *Trans. Am. Math. Soc.* **295**, 607–633 (1986)
21. Preston, G.B.: Inverse semigroups. *J. Lond. Math. Soc.* **29**, 396–403 (1954)
22. Reither, S.: *Die natürliche ordnung auf halbgruppen*. Ph.D. Thesis, University of Vienna (1994)
23. Sen, M.K.: \mathcal{P} -congruences on \mathcal{P} -regular semigroups. *Semigroup Forum* **44**, 149–156 (1992)
24. Siripitukdet, M., Sattayaporn, S.: The maximum idempotent-separating congruence on E -inversive E -semigroups. *Int. J. Algebra* **2**, 925–931 (2008)
25. Thierrin, G.: Sur les demi-groupes inverses. *C. R. Acad. Sci. Paris* **234**, 1336–1338 (1952)
26. Trotter, P.G.: Normal partitions of idempotents of regular semigroups. *J. Aust. Math. Soc. A* **26**, 110–114 (1978)
27. Weipoltshammer, B.: Certain congruences on E -inversive E -semigroups. *Semigroup Forum* **65**, 233–248 (2002)
28. Yamada, M., Sen, M.K.: \mathcal{P} -regular semigroups. *Semigroup Forum* **39**, 157–178 (1989)
29. Zheng, H.W.: Strong \mathcal{P} -congruences on \mathcal{P} -regular semigroups. *Semigroup Forum* **51**, 217–223 (1995)
30. Zheng, H.W.: Group congruences on an E -inversive semigroup. *Southeast Asian Bull. Math.* **21**, 1–8 (1997)